

Determination of phylogroup in extended spectrum beta lactamase (ESBL)-*E. coli* from fishes by Clermont's rapid phylotyping method

Sivaraman G.K.^{1*}, Sudha S.¹, Muneeb K.H.¹, Jennifer Cole²,
Bibek R. Shome³ and Mark Holmes⁴

¹ICAR-Central Institute of Fisheries Technology, Cochin - 29

²Department of Geography, Royal Holloway, University of London, UK.

³ICAR-National Institute of Veterinary Epidemiology and Disease Informatics, Bangalore

⁴Department of Veterinary Medicine, University of Cambridge, UK.

*gkshivraman@gmail.com

Escherichia coli is a normal intestinal inhabitant in majority of the animals including humans. It is a versatile commensal but capable of causing both intestinal and extraintestinal diseases such as diarrhoea, septicaemia, urinary tract infections and neonatal meningitis (Orskov and Orskov, 1992). The strains of *E. coli* are diverse with its genome plasticity and the population structure of this organism remains mostly clonal (Touchon et al., 2009). Despite this, *E. coli* can be delineated into six prominent phylogenetic groups-A, B1, B2, D, E and F. Determining the phylogroup of *E. coli* help to understand the path-ogenicity and also give an insight of how these pathogenic strains acquire virulence genes. Tech-niques like multi locus enzyme electrophoresis and ribotyping are capable of assigning phylogroups, but these are time consuming, complex and need a collection of typed strains (Clermont et al., 2000). In 2000, Clermont et al., (2000) developed a PCR based method to characterize the phylogroup based on three candidate markers

—*chuA*, *yjaA* and *TSPE4.C2*. The primers designed were such that strains could be classified into four groups A, B1, B2 and D. Extensive multi locus sequence data and genome data of *E. coli* lead the Clermont group to further improvise the method and in 2013, characterization of eight phylogroups (A, B1, B2, C, D, E, F and *E. coli* cryptic clade I) of *E. coli* were made possible.

The recent study documented 50 extended spectrum beta lactamase (ESBL) *E. coli* isolated and characterized from fishes collected from Guwahati, Assam through quadraplex PCR phylotyping method of Clermont et al., (2013) (Sivaraman et al., 2020). *arpA* gene was used as an internal control for DNA quality and the *E. coli* and *Escherichia* clade I strains were expected to give a positive amplicon. Other than the primers used in the quadraplex, two other primers viz., *trpAggC* and *ArpAggE* (Lescat et al., 2012) were also used to classify phylogroups C and E, respectively. *trp BA* was used to provide an internal control. The primer sequences and the PCR conditions are listed in Table 1.

Table1: Primer sequence, reaction conditions and amplicon size of the gene targets

Target	Primer Name	Sequence (5'-3')	Length (bases)	Primer concentration (μM)	Annealing temp. (°C)	Amplicon size (bp)
<i>chuA</i>	<i>chuA.1b</i>	ATGGTACCGGACGAACCAAC	20	1.0	59	288
	<i>chuA.2</i>	TGCCGCCAGTACCAAAGACA	20	1.0		
<i>yjaA</i>	<i>yjaA.1b</i>	CAAACGTGAAGTGTCAGGAG	20	1.0		211
	<i>yjaA.2b</i>	AATGCGTTCCTCAACCTGTG	20	1.0		
<i>TspE4.C2</i>	<i>TspE.C2.1b</i>	CACTATTCGTAAGGTCATCC	20	1.0		152
	<i>TspE4C2.2b</i>	AGTTTATCGCTGCGGGTCGC	20	1.0		

<i>arpA</i>	<i>AceK.f</i> <i>ArpA1.r</i>	AACGCTATTCGCCAGCTTGC TCTCCCCATACCGTACGCTA	20 20	2.0 2.0		400
<i>arpA</i> – Group E	<i>ArpAgpE.f</i> <i>ArpAgpE.r</i>	GATTCCATCTTGTCAAAATATGCC GAAAAGAAAAAGAATTCCCAAGAG	24 24	1.0 1.0	57	301
<i>trpA</i> - Group C	<i>trpAgpC.1</i> <i>trpAgpC.2</i>	AGTTTATGCCCAGTGCGAG TCTGCGCCGGTCAACGCC	20 18	1.0 1.0	59	219
<i>trpA</i> -Internal control	<i>trpBA.f</i> <i>trpBA.r</i>	CGGCGATAAAGACATCTTCAC GCAACGCGGCCTGGCGGAAG	21 20	0.6 0.6	57/59	489

The results showed that forty percent ($n=20$) of the *E. coli* strains were grouped into phylogroup B1 and 30% ($n=15$) to group A. Other prominent phylogroups amongst the strains were E (12%, 6) and D (8%, 4). Two strains belonged to group F and one to Group C. One isolate could not be typed based on Clermont phylogrouping and was termed 'Unknown' as the PCR results were negative for *arpA/chuA/yjaA/TspE4.C2*. This isolate was positive when screened for *E. coli* specific *uidA* gene. The Fig 1 shows the amplification results of few strains studied. Literature points to the fact that virulent extra-intestinal strains of *E. coli* mainly belong to phylogroup B2 and to a smaller extent to Group D. Most of the commensal strains belong to phylogroup A. Phylogroup E was earlier unassigned to which one of the most important serotype O157:H7 belongs and group F is a sister group of phylogroup B2. Group C is closely related to but distinct from phylogroup B1. These findings emphasizes the phylogenetic diversity of *E. coli* strains isolated from fish source and their ability to disseminate infection. In humans and animals, intra-intestinal infections are caused mostly by phylogroup A/B1 or E and extra-intestinal infections are caused mainly by strains belonging to B2 (Clermont et al., 2011). The findings are in concordance with the

reports that *E. coli* strains from animals usually belong to phylogroup B1 with 34-50% incidence. None of the studied strains belonged to pathogenic B2 phylogroup. About 8% of the *E. coli* strains belonged to phylogroup D which is known to be present in extra-intestinal pathogenic strains and calls for greater alert. The study validates Clermont's phylotyping method as a simple, rapid, robust and inexpensive tool for phylogrouping of *E. coli*.

Funding

This study was supported by Department of Biotechnology (DBT), Government of India (BT/IN/indo-UK/AMR/06/BRS/2018-19) and Economic and Social Research Council, UK.




Fig1 : Quadraplex PCR with the primer combination -*arpA/chuA/yjaA/TspE4.C2*.

References:

- Ørskov, F., & Ørskov, I. (1992). *Escherichia coli* serotyping and disease in man and animals. *Canadian journal of microbiology*, 38(7), 699-704.
- Touchon, M., Hoede, C., Tenaillon, O., Barbe, V., Baeriswyl, S., Bidet, P., ... & Denamur, E. (2009). Organised genome dynamics in the *Escherichia coli* species results in highly diverse adaptive paths. *PLoS genet*, 5(1), e1000344.
- Clermont, O., Bonacorsi, S., & Bingen, E. (2000). Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Applied and environmental microbiology*, 66(10), 4555-4558.
- Sivaraman, G. K., Sudha, S., Muneeb, K. H., Shome, B., Holmes, M., & Cole, J. (2020). Molecular assessment of antimicrobial resistance and virulence in multi drug resistant ESBL-producing *Escherichia coli* and *Klebsiella pneumoniae* from food fishes, Assam, India. *Microbial Pathogenesis*, 149, 104581.
- Clermont, O., Christenson, J. K., Denamur, E., & Gordon, D. M. (2013). The Clermont *Escherichia coli* phylo-typing method revisited: improvement of specificity and detection of new phylo-groups. *Environmental microbiology reports*, 5(1), 58-65.
- Lescat, M., Clermont, O., Woerther, P. L., Glodt, J., Dion, S., Skurnik, D. & Denamur, E. (2013). Commensal *Escherichia coli* strains in Guyana reveal a high genetic diversity with host dependant population structure. *Environmental microbiology reports*, 5(1), 49-57.
- Clermont, O., Olier, M., Hoede, C., Diancourt, L., Brisse, S., Keroudean, M., Glodt, J., Picard, B., Oswald, E., & Denamur, E. (2011). Animal and human pathogenic *Escherichia coli* strains share common genetic backgrounds. *Infection, Genetics and Evolution : Journal of molecular epidemiology and evolutionary genetics in infectious diseases*, 11(3), 654-662.